Asian mussel watch program: Contamination status of polybrominated diphenyl ethers and organochlorines in coastal waters of Asian countries

Ramu K., Kajiwara N., Sudaryanto A., Isobe T., Takahashi S., Subramanian A., Ueno D., Zheng G.J., Lam P.K.S., Takada H., Zakaria M.P., Viet P.H., Prudente M., Tana T.S., Tanabe S.

Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan; Division of Environmental Conservation, Saga University, Japan; Department of Biology and Chemistry, City University of Hong Kong, Hong Kong, Hong Kong; Laboratory of Organic Geochemistry, Tokyo University of Agriculture and Technology, Japan; Faculty of Environmental Studies, Universiti Putra Malaysia, Malaysia; Center for Environmental Technology and Sustainable Development, Hanoi National University, Viet Nam; Science Education Department, De La Sella University, Philippines; Social and Cultural Observation Unit (OBSES), Office of the Council of Ministers, Cambodia

Abstract: Mussel samples were used in this study to measure the levels of polybrominated diphenyl ethers (PBDEs) and organochlorines (OCs) in the coastal waters of Asian countries like Cambodia, China, Hong Kong, India, Indonesia, Japan, Korea, Malaysia, the Philippines, and Vietnam. PBDEs were detected in all the samples analyzed, and the concentrations ranged from 0.66 to 440 ng/g lipid wt. Apparently higher concentrations of PBDEs were found in mussels from the coastal waters of Korea, Hong Kong, China, and the Philippines, which suggests that significant sources of these chemicals exist in and around this region. With regard to the composition of PBDE congeners, BDE-47, BDE-99, and BDE-100 were the dominant congeners in most of the samples. Among the OCs analyzed, concentrations of DDTs were the highest followed by PCBs > CHLs > HCHs > HCB. Total concentrations of DDTs, PCBs, CHLs, and HCHs in mussel samples ranged from 21 to 58 000, 3.8 to 2000, 0.93 to 900, and 0.90 to 230 ng/g lipid wt., respectively. High levels of DDTs were found in mussels from Hong Kong, Vietnam, and China; PCBs were found in Japan, Hong Kong, and industrialized/urbanized locations in Korea, Indonesia, the Philippines, and India; CHLs were found in Japan and Hong Kong; HCHs were found in India and China. These countries seem to play a role as probable emission sources of corresponding contaminants in Asia and, in turn, may influence their global distribution. © 2007 American Chemical Society.

Index Keywords: Coastal waters; Polybrominated diphenyl ethers (PBDE); Coastal zones; Concentration (process); Contamination; Ethers; Fish; Water pollution; organochlorine derivative; polybrominated diphenyl ether; sea water; Coastal zones; Concentration (process); Contamination; Ethers; Fish; Water pollution; bioaccumulation; bioindicator; biomonitoring; bivalve; coastal water; DDT; emission inventory; geographical distribution; organochlorine; PBDE; PCB; pollutant source; water pollution; article; Asia; Cambodia; China; coastal waters; comparative study; geographic distribution; Hong Kong; India; Indonesia; industrial area; Japan; Korea; Malaysia; mussel; nonhuman; Philippines; sea pollution; Viet Nam; water pollutant; Animals; Asia; Bivalvia; Geography; Hydrocarbons, Chlorinated; Phenyl Ethers; Polybrominated Biphenyls; Water Pollutants, Chemical; Asia; Cambodia; China; Eurasia; Far East; Hong Kong; India; Indonesia; Japan; Korea; Malaysia; Philippines; South Asia; Southeast Asia; Viet Nam

Year: 2007

Source title: Environmental Science and Technology

Volume: 41 Issue: 13

Page: 4580-4586

Cited by: 26

Link: Scorpus Link

Chemicals/CAS: Hydrocarbons, Chlorinated; Phenyl Ethers; Polybrominated Biphenyls; Water Pollutants, Chemical; pentabromodiphenyl ether, 32534-81-9

Correspondence Address: Tanabe, S.; Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan; email: shinsuke@agr.ehime-u.ac.jp

ISSN: 0013936X CODEN: ESTHA

DOI: 10.1021/es070380p PubMed ID: 17695900

Language of Original Document: English

Abbreviated Source Title: Environmental Science and Technology

Document Type: Article

Source: Scopus

Authors with affiliations:

- Ramu, K., Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan
- Kajiwara, N., Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan
- Sudaryanto, A., Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan
- Isobe, T., Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan
- Takahashi, S., Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan
- Subramanian, A., Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan
- Ueno, D., Division of Environmental Conservation, Saga University, Japan
- Zheng, G.J., Department of Biology and Chemistry, City University of Hong Kong, Hong Kong, Hong Kong
- Lam, P.K.S., Department of Biology and Chemistry, City University of Hong Kong, Hong Kong, Hong Kong
- Takada, H., Laboratory of Organic Geochemistry, Tokyo University of Agriculture and Technology, Japan
- Zakaria, M.P., Faculty of Environmental Studies, Universiti Putra Malaysia, Malaysia
- Viet, P.H., Center for Environmental Technology and Sustainable Development, Hanoi National University, Viet Nam
- Prudente, M., Science Education Department, De La Sella University, Philippines
- Tana, T.S., Social and Cultural Observation Unit (OBSES), Office of the Council of Ministers, Cambodia
- Tanabe, S., Center for Marine Environmental Studies, Ehime University, Bunkyo-cho 2-5, Matsuyama 790-8577, Japan References:
- Hites, R.A., Polybrominated diphenyl ethers in the environment and in people: A meta-analysis of concentrations (2004) Environ. Sci. Technol, 38, pp. 945-956
- Hileman, B., Electronic waste (2002) Chem. Eng. News, 80 (26), pp. 15-18
- Hale, R.C., Guardia, M.J.L., Harvey, E., Mainor, T.M., Potential role of fire retardant-treated polyurethane foam as a source of brominated diphenyl ethers to the US environment (2002) Chemosphere, 46, pp. 729-735
- Dave, P.P.I., A generic giant (1996) Farm. Chem. Int, 10, pp. 36-37

- Li, Y.F., Mcmillan, A., Scholtz, M.T., Global HCH usage with 1° x 1° longitude/latitude resolution (1996) Environ. Sci. Technol, 30, pp. 3525-3533
- Monirith, I., Ueno, D., Takahashi, S., Nakata, H., Sudaryanto, A., Subramanian, A., Karuppiah, S., Tanabe, S., Asia-Pacific mussel watch: Monitoring contamination of persistent organochlorine compounds in coastal waters of Asian countries (2003)
 Mar. Pollut. Bull, 46, pp. 281-300
- Ueno, D., Takahashi, S., Tanaka, H., Subramanian, A., Fillman, G., Nakata, H., Lam, P.K.S., Tanabe, S., Global pollution monitoring of PCBs and organochlorine pesticides using skipjack tuna as a bioindicator (2003) Arch. Environ. Contam. Toxicol, 45, pp. 378-389
- Jeng, M.-S.
- Jeng, W.-L.
- Hung, T.-C.
- Yeh, C.-Y.
- Tseng, R.-J.
- Meng, P.-J.
- Han, B.-C. Mussel Watch: a review of Cu and other metals in various marine organisms in Taiwan, 1991-98. Environ. Pollut. 2000, 110, 207-2150', Connor, T. P.
- Lauenstein, G. G. Trends in chemical concentrations in mussels and oysters collected along the US coast: Update to 2003. Mar. Environ. Res. 2006, 62, 261-285Isobe, T., Takada, H., Kanai, M., Tsutsumi, S., Isobe, K.O., Boonyatumanond, R., Zakaria, M.P., Distribution of polycyclic aromatic hydrocarbons (PAHs) and phenolic endocrine disrupting chemicals in South and Southeast Asian mussels (2007) Environ. Monit. Assess, , in press
- Ramu, K., Kajiwara, N., Isobe, T., Takahashi, S., Kim, E.-Y., Min, B.-Y., We, S.-U., Tanabe, S., Spatial distribution and accumulation of brominated flame retardants, polychlorinated biphenyls and organochlorine pesticides in blue mussels (Mytilus edulis) from coastal waters of Korea (2007) Environ. Pollut, , in press
- Ueno, D., Kajiwara, N., Tanaka, H., Subramanian, A., Fillmann, G., Lam, P.K.S., Zheng, G.J., Tanabe, S., Global pollution monitoring of polybrominated diphenyl ethers using skipjack tuna as a bioindicator (2004) Environ. Sci. Technol, 38, pp. 2312-2316
- Kajiwara, N., Ueno, D., Monirith, I., Tanabe, S., Pourkazemi, M., Aubrey, D.G., Contamination by organochlorine compounds in sturgeons from Caspian Sea during 2001 and 2002 (2003) Mar. Pollut. Bull, 46, pp. 741-747
- Johansson, I., Moisan, K.H., Guiot, N., Munschy, C., Tronczyński, J., Polybrominated diphenyl ethers (PBDEs) in mussels from selected French coastal sites: 1981-2003 (2006) Chemosphere, 64, pp. 296-305
- Oros, D.R., Hoover, D., Rodigari, F., Crane, D., Sericano, J., Levels and distribution of polybrominated diphenyl ethers in water, surface sediments and bivalves from the San Francisco Estuary (2005) Environ. Sci. Technol, 39, pp. 33-41
- Alaee, M., Arias, P., Sjodin, A., Bergman, A., An overview of commercially used brominated flame retardants, their applications, their use patterns in different countries/regions and possible modes of release (2003) Environ. Int, 29, pp. 683-689
- Stapleton, H.M., Alaee, M., Letcher, R.J., Baker, J.E., Debromination of the flame retardant decabromodiphenyl ether by juvenile carp (Cyprinus carpio) following dietary exposure (2004) Environ. Sci. Technol, 38, pp. 112-119
- Mai, B., Chen, S., Luo, X., Chen, L., Yang, Q., Sheng, G., Peng, P., Zeng, E.Y., Distribution of polybrominated diphenyl ethers in sediments of the Pearl River Delta and adjacent South China Sea (2005) Environ. Sci. Technol, 39, pp. 3521-3527
- Moon, H.-B., Kannan, K., Lee, S.-J., Choi, M., Polybrominated diphenyl ethers (PBDEs) in sediment and bivalves from Korean coastal waters (2007) Chemosphere, 66, pp. 243-251
- de Boer, J., Wester, P.G., Horst, A.V., Leonards, P.E.G., Polybrominated diphenyl ethers in influents, suspended particulate

- matter, sediments, sewage treatment plant and effluents and biota from the Netherlands (2003) Environ. Pollut, 122, pp. 63-74
- de Wit, C.Y., An overview of brominated flame retardants in the environment (2002) Chemosphere, 46, pp. 583-624
- (2004), http://www.bsef.com, Bromine Science and Environmental Forum BSEFWang, X.M., Ding, X., Mai, B.X., Xie, Z.Q., Xiang, C.H., Sun, L.G., Sheng, G.Y., Zeng, E.Y., Polybrominated diphenyl ethers in airborne particulates collected during a research expedition from the Bohai Sea to the Arctic (2005) Environ. Sci. Technol, 39, pp. 7803-7809
- Zheng, G.J., Martin, M., Richardson, B.J., Yu, H., Liu, Y., Zhou, C., Li, J., Lam, P.K.S., Concentrations of polybrominated diphenyl ethers (PBDEs) in Pearl River Delta sediments (2004) Mar. Pollut. Bull, 49, pp. 520-524
- Christensen, J.H., Platz, J., Screening of polybrominated diphenyl ethers in blue mussels, marine and freshwater sediments in Denmark (2001) J. Environ. Monit, 3, pp. 543-547
- Leung, A.O.W., Luksemburg, W.J., Wong, A.S., Wong, M.H., Spatial distribution of polybrominated diphenyl ethers and polychlorinated dibenzo-p-dioxins and dibenzofurans in soil and combusted residue at Guiyu, an electronic waste recycling site in Southeast China Environ. Sci. Technol, 41, pp. 2730-2737
- Liu, Y., Zheng, G.J., Yu, H., Martin, M., Richardson, B.J., Lam, M.H.W., Lam, P.K.S., Polybrominated diphenyl ethers (PBDEs) in sediments and mussel tissues from Hong Kong marine waters (2005) Mar. Pollut. Bull, 50, pp. 1173-1184
- Hale, R.C., LaGuardia, M.J., Harvey, E.P., Gaylor, M., Matteson, M.Y., Duff, W.H., Persistent pollutants in land applied sludges (2001) Nature, 412, pp. 140-141
- Ramu, K., Kajiwara, N., Tanabe, S., Lam, P.K.S., Jefferson, T.A., Polybrominated diphenyl ethers (PBDEs) and organochlorines in small cetaceans from Hong Kong waters: Levels, profiles and distribution (2005) Mar. Pollut. Bull, 51, pp. 669-676
- Klumpp, D. W.
- · Huasheng, H.
- · Humphrey, C.
- · Wang, Xinhong, W.
- Codia, S. Toxic contaminants and their biological effects in coastal waters of Xiamen, China. I. Organic pollutants in mussel and fish tissues. Mar. Pollut. Bull. 2002, 44, 752-760Qiu, X., Zhu, T., Yao, B., Hu, J., Hu, S., Contribution of dicofol to the current DDT pollution in China (2005) Environ. Sci. Technol, 39, pp. 4385-4390
- Wu, Y., Zhang, J., Zhou, Q., Persistent organic residues in sediments from Chinese river/estuary systems (1999) Environ. Pollut, 105, pp. 143-150
- Yuan, D., Yang, D., Wade, T.L., Qian, Y., Status of persistent organic pollutants in the sediment from several estuaries in China (2001) Environ. Pollut, 114, pp. 101-111
- Jaward, F.M., Zhang, G., Nam, J.J., Sweetman, A.J., Obbard, J.P., Kobara, Y., Jones, K.C., Passive air sampling of polychlorinated biphenyls, organochlorine compounds and polybrominated diphenyl ethers across Asia (2005) Environ. Sci. Technol, 39, pp. 8638-8645
- Kajiwara, N., Kamikawa, S., Ramu, K., Ueno, D., Yamada, T.K., Subramanian, A., Lam, P.K.S., Tanabe, S., Geographical distribution of polybrominated diphenyl ethers (PBDEs) and organochlorines in small cetaceans from Asian waters (2006) Chemosphere, 64, pp. 287-295
- Loganathan, B.G., Tanabe, S., Hidaka, Y., Kawano, M., Hidaka, H., Tatsukawa, R., Temporal trends of persistent organochlorine residues in human adipose tissue from Japan, 1928-1985 (1993) Environ. Pollut, 81, pp. 31-39
- Fu, J., Mai, B., Sheng, G., Zhang, G., Wang, X., Peng, P., Xiao, X., Tang, U.W., Persistent organic pollutants in environment of the Pearl River Delta, China: An overview (2003) Chemosphere, 52, pp. 1411-1422
- Li, Y.F., Cai, D.J., Singh, A., Technical hexachlorocyclohexane use trends in China and their impact on the environment (1998)

Arch. Environ. Contam. Toxicol, 35, pp. 668-697

- Li, Y.F., Global technical hexachlorocyclohexane usage and its contamination consequences in the environment: From 1948 to 1997 (1999) Sci. Total Environ, 232, pp. 121-158
- Xu, D., Deng, L., Chai, Z., Mao, X., Organohalogenated compounds in pine needles from Beijing city, China (2004) Chemosphere, 57, pp. 1343-1353
- Pandit, G.G., Sahu, S.K., Sharma, S., Puranik, V.D., Distribution and fate of persistent organochlorine pesticides in coastal marine environment of Mumbai (2006) Environ. Int, 32, pp. 240-243
- Wania, F., Mackay, D., Tracking the distribution of persistent organic pollutants (1996) Environ. Sci. Technol, 30, pp. 390A-396A